

New prospects for PCSK9 inhibition?

Ulf Landmesser^{1*}, M. John Chapman², Jane K. Stock³, Pierre Amarenco⁴, Jill J. F. Belch⁵, Jan Borén⁶, Michel Farnier⁷, Brian A. Ference⁸, Stephan Gielen⁹, Ian Graham¹⁰, Diederick E. Grobbee¹¹, G. Kees Hovingh¹², Thomas F. Lüscher¹³, Massimo F. Piepoli¹⁴, Kausik K. Ray¹⁵, Erik S. Stroes¹², Olov Wiklund¹⁶, Stephan Windecker¹⁷, Jose Luis Zamorano¹⁸, Fausto Pinto¹⁹, Lale Tokgözoğlu²⁰, Jeroen J. Bax²¹, and Alberico L. Catapano²²; European Society of Cardiology/
 European Atherosclerosis Society Task Force

¹Department of Cardiology, Charité - Universitätsmedizin Berlin (CBF), and Institute of Health (BIH), Berlin, Germany; ²National Institute for Health and Medical Research (INSERM), University of Pierre and Marie Curie, Pitié-Salpêtrière Hospital, Paris, France; ³European Atherosclerosis Society, Gothenburg, Sweden; ⁴Department of Neurology and Stroke Centre, Bichat Hospital, Paris-Diderot-Sorbonne University, Paris, France; ⁵Institute of Cardiovascular Research, Ninewells Hospital and Medical School, Dundee, UK; ⁶Department of Molecular and Clinical Medicine, University of Gothenburg and Sahlgrenska University Hospital, and Wallenberg Laboratory, Sahlgrenska University Hospital, Gothenburg, Sweden; ⁷Lipid Clinic, Point Medical, and Department of Cardiology, CHU Dijon-Bourgogne, Dijon, France; ⁸Division of Cardiovascular Medicine, Division of Translational Research and Clinical Epidemiology, Wayne State University School of Medicine, Detroit, MI, USA; ⁹Martin-Luther-University Halle/Wittenberg, University Hospital, Department of Internal Medicine III, Halle/Saale, Germany; ¹⁰Trinity College Dublin, Ireland; ¹¹Julius Global Health, the Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, The Netherlands; ¹²Academic Medical Center, Department of Vascular Medicine, University of Amsterdam, Amsterdam, The Netherlands; ¹³University Heart Center, Department of Cardiology, University Hospital Zurich, and Center for Molecular Cardiology, University of Zurich, Zurich, Switzerland; ¹⁴G Da Saliceto Hospital, Heart Failure Unit, Cardiac Department, Piacenza, Italy; ¹⁵Imperial Centre for Cardiovascular Disease Prevention, Department of Primary Care and Public Health, Imperial College, London, UK; ¹⁶Sahlgrenska University Hospital, Gothenburg, Sweden; ¹⁷Department of Cardiology, Swiss Cardiovascular Center, University Hospital, Bern, Switzerland; ¹⁸Department of Cardiology, University Hospital Ramón y Cajal, Madrid, Spain; ¹⁹Cardiology Department, CCUL, CAML, Faculdade de Medicina, Universidade de Lisboa, Portugal; ²⁰Department of Cardiology, Hacettepe University, Ankara, Turkey; ²¹Department of Cardiology, Leiden University Medical Center, Leiden, The Netherlands; and ²²University of Milan and Multimedica IRCSS Milano, Italy

Online publish-ahead-of-print 22 March 2018

This commentary refers to ‘2017 update of ESC/EAS task force on practical clinical guidance for proprotein convertase subtilisin/kexin type 9 inhibition in patients with atherosclerotic cardiovascular disease or in familial hypercholesterolaemia’, by U Landmesser et al., 2018;39: 1131–1143.

In 2017, this Task Force updated practical guidance for clinical use of proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibition following publication of FOURIER.¹ Beyond licenced indications, PCSK9 inhibitors may have application in other high-risk conditions, such as severe hyperlipidaemia with liver failure,² supported by pharmacodynamic data,³ although the lack of trials to date does not allow recommendations.

Recent insights from FOURIER help to define patients at highest risk with elevated low-density lipoprotein cholesterol (LDL-C) levels who benefit most from PCSK9 inhibition. These include those with symptomatic peripheral artery disease (PAD), a group often under-recognized and undertreated, in whom evolocumab reduced major adverse cardiovascular events (MACE) by 27% and major adverse limb events by 37%, with benefits extending to LDL-C levels <0.26 mmol/L.⁴ Patients with recent or recurrent myocardial

infarction (MI), or multivessel disease, at 34–90% higher risk of a MACE, also derived greater benefit from evolocumab than those without these characteristics (*Sabatine MS, Annual Scientific Sessions, American Heart Association, 13 November 2017*). Thus, irrespective of other vascular disease, symptomatic PAD, the timing and frequency of MI, or multivessel disease, associated with residual LDL-C burden, indicate very high-risk patients who merit consideration of PCSK9 inhibition.

Will results from ODYSSEY OUTCOMES extend use of PCSK9 inhibitors to early post-MI patients? Such findings would align with the MIRACL trial⁵ and may imply further stabilization of atherosclerotic plaque from LDL-C lowering with PCSK9 inhibition.

Conflict of interest: Potential conflicts of interest of the authors are denoted in reference 1.

References

- Landmesser U, Chapman MJ, Stock JK, Amarenco P, Belch JFF, Borén J, Farnier M, Ference BA, Gielen S, Graham I, Grobbee DE, Hovingh GK, Lüscher TF, Piepoli MF, Ray KK, Stroes ES, Wiklund O, Windecker S, Zamorano JL, Pinto F, Tokgözoğlu L, Bax JJ, Catapano AL. 2017 update of ESC/EAS task force on practical clinical guidance for proprotein convertase subtilisin/kexin type 9 inhibition in patients with atherosclerotic cardiovascular disease or in familial hypercholesterolaemia. *Eur Heart J* 2018;**39**:1131–1143.

* Corresponding author. Email: ulf.landmesser@charite.de

Published on behalf of the European Society of Cardiology. All rights reserved. © The Author(s) 2018. For permissions, please email: journals.permissions@oup.com.

2. Glowczynska R, Hus A, Raszeja-Wyszomirska J, Opolski G. Are the proprotein convertase subtilisin/kexin type 9 inhibitors new therapeutic chance for patients with familial hypercholesterolaemia and liver failure? *Eur Heart J* 2018; doi: 10.1093/eurheartj/ehy361.
3. Gibbs JP, Slatter JG, Egbuna O, Geller M, Hamilton L, Dias CS, Xu RY, Johnson J, Wasserman SM, Emery MG. Evaluation of evolocumab (AMG 145), a fully human anti-PCSK9 IgG2 monoclonal antibody, in subjects with hepatic impairment. *J Clin Pharmacol* 2017;**57**:513–523.
4. Bonaca MP, Nault P, Giugliano RP, Keech AC, Pineda AL, Kanevsky E, Kuder J, Murphy SA, Jukema JW, Lewis BS, Tokgozoglu L, Somaratne R, Sever PS, Pedersen TR, Sabatine MS. Low-density lipoprotein cholesterol lowering with evolocumab and outcomes in patients with peripheral artery disease: insights from the FOURIER trial (Further Cardiovascular Outcomes Research With PCSK9 Inhibition in Subjects With Elevated Risk). *Circulation* 2018;**137**: 338–350.
5. Schwartz GG, Olsson AG, Ezekowitz MD, Ganz P, Oliver MF, Waters D, Zeiher A, Chaitman BR, Leslie S, Stern T. Myocardial Ischemia Reduction with Aggressive Cholesterol Lowering (MIRACL) Study Investigators. Effects of atorvastatin on early recurrent ischemic events in acute coronary syndromes: the MIRACL study: a randomized controlled trial. *Jama* 2001;**285**:1711–1718.